





ONR-TR-21 AD-AO88	3. RECIPIENT'S CATALOG NUMBER					
TITLE (and Substitio)	Y AX A					
TITLE (und Subtitio)	QQ0 [9]					
Dhatachamietus of Cabalt Caubanul Compleyee Mauin	S. TYPE OF REPORT & PERIOD COVERE					
Photochemistry of Cobalt Carbonyl Complexes Having a Cobalt-Silicon Bond and its Importance in	Inter in recinical kepar					
Activation of Catalysis.	6. PERFORMING ORG. REPORT NUMBER					
AUTHOR(a)	8. CONTRACT OF GRANT NUMBER(*) NØ6014-75-C-Ø88Ø					
Carol L. Reichel and Mark S. Wrighton	1100014-73-0-1000					
Department of Chemistry	16. PROGRAM ELEMENT, PROJECT, TAS AREA & WORK UNIT NUMBERS					
Massachusetts Institute of Technology	NR-051-579					
Cambridge, MA 02139	NK-U51-5/9					
1. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE					
Office of Naval Research 23 70 80	August 22, 1980					
	22					
Arlington VA 22217  4. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)	18. SECURITY CLASS. (of this report) UNCLASSIFIED					
$(P, g_3)$	UNCLASSIFIED					
	184. DECLASSIFICATION/DOWNGRADING					
16. DISTRIBUTION STATEMENT (of this Report)						
Approved for public release; reproduction is pe of the United States Government; distribution unl	DTIC ELECTE					
7. DISTRIBUTION STATEMENT (of the ebetrect entered in Block 20, if different is Distribution of this document is unlimited.	SEP 3 1980					
8. SUPPLEMENTARY NOTES	D					
Prepared and accepted for publication in <u>Inorganic Chemistry</u> .						
9. KEY WORDS (Continue on severee side if necessary and identify by block number	n)					

201 ABSTRACT (Continue on reverse side if necessary and identify by block number)

1+01-

Photochemistry of  $R_3^{\prime\prime}SiCo(CO)_4^{\prime\prime}$  (R = Et, Ph) in hydrocarbon solvents shows that the dominant primary excited state process is loss of CO, not cleavage of the Si-Co bond. The 366 nm quantum yield for CO substitution by  $P(OPh)_3$  is  $0.28 \pm 0.05$  to give  $R_3SiCo(CO)_3P(OPh)_3$ . Irradiation of  $R_3SiCo(CO)_4$  in the presence of  $0.1 \, \text{M} \, \text{HSiR'}_3$  gives  $R'_3SiCo(CO)_4$  with a quantum yield nearly the same as that for CO substitution by  $P(OPh)_3$ ; irradiation of  $Ph_3SiCo(CO)_4$  in the presence of  $0.1 \, \text{M} \, \text{HSiEt}_3$  and  $0.3 \, \text{M} \, P(OPh)_3$  yields only  $Ph_3SiCo(CO)_3P(OPh)_3$ , not

DD 1 JAN 73 1473

hydrosilation.

UNCLASSIFIED

(over)

UNCLASSIFIED

HTY CLASSIFICATION OF THIS PAGE(When Date Entered)  $(Co(CO)_3P(OPh)_3$ . Irradiation of  $R_3SiCo(CO)_4$  in the presence of 1-pentene First gives RaSiCo(CO) (alkene), but the dominant Co-containing material is ultimately a mixture  $Co_2(CO)_8$  and  $Co_4(CO)_{12}$  at 25°C. For R=Et, the dominant Si-containing products have been determined to be isomers of (pentenyl)SiEt3. The quantum yield for reaction of Et<sub>3</sub>SiCo(CO)<sub>4</sub> with 1-pentene ( $\Phi$  = 0.23) is significantly higher than the reaction quantum yield for Et<sub>3</sub>SiCo(CO)<sub>4</sub> in isooctane ( $\Phi = 0.04$ ) solution where mixtures of  $Co_2(CO)_8$  and  $Co_4(CO)_{12}$  are the photoproducts. Irradiation of R₃SiCo(CO), in the presence of 1-pentene/HSiEt₃ (1/1 mole ratio) yields catalytic alkene isomerization and hydrosilation. Thermal catalysis (25°C) of 1-pentene hydrosilation by HSiEt<sub>3</sub> using Co<sub>2</sub>(CO)<sub>8</sub> yields infrared detectable Et<sub>3</sub>SiCo(CO)<sub>4</sub>. The Et<sub>3</sub>SiCo(CO)<sub>4</sub> does not catalyze thermal reaction at 25°C at a rate comparable to that initially observed using Co<sub>2</sub>(CO)<sub>8</sub>. Regeneration of catalytic activity in the Co<sub>2</sub>(CO)<sub>8</sub>/1-pentene/HSiEt<sub>3</sub> system can be accomplished photochemically by exciting the Et'SiCo(CO), species.

1	. ៤៤៩ន	ion For	
!	l IS D. TA	В	A
	Unamno Justii	unced ication_	
	Ву		
	Distri	bution/	
Ì	Avail	ability	Codes
	Dist	Avail an specia	

OFFICE OF NAVAL RESEARCH

Contract NO0014-75-C-0880

Task No. NR 051-579

TECHNICAL REPORT NO. 21

# PHOTOCHEMISTRY OF COBALT CARBONYL COMPLEXES HAVING A COBALT-SILICON BOND AND ITS IMPORTANCE IN ACTIVATION OF CATALYSIS

bу

Carol L. Reichel and Mark S. Wrighton Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139

Prepared for Publication
in
INORGANIC CHEMISTRY

August 22, 1980

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

D

Contribution from Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Photochemistry of Cobalt Carbonyl Complexes Having a Cobalt-Silicon Bond and Its Importance in Activation of Catalysis

Carol L. Reichel and Mark S. Wrighton\*

<sup>\*</sup>To whom to address correspondence should be addressed.

We wish to report results of experiments that establish the relative importance of photochemical cleavage of the Co-Si bond vs. the dissociative loss of CO in  $R_3SiCo(CO)_4$ , equation (1) vs. (2). We recently showed that


$$R_3 \text{SiCo(CO)}_4 \xrightarrow{hv} R_3 \text{Si} + \text{Co(CO)}_4$$
 (1)

$$R_3 Sico(CO)_4 \xrightarrow{hv} R_3 Sico(CO)_3 + CO$$
 (2)

photogeneration of 17-valence electron, Co-centered radicals can lead to catalysis of alkene isomerization, hydrosilation, and hydrogenation. Reaction of  $R_3 SiCo(CO)_4$  according to equation (1) is of interest in that the 17-valence electron radical  $\cdot Co(CO)_4$  could effect the same alkene/HSiR $_3$  reactions known $^2$  to be catalyzed by  $Co_2(CO)_8$ . It is believed that  $R_3 SiCo(CO)_4$  represents the ultimate fate of Co in the  $Co_2(CO)_8$  catalyzed hydrosilation reaction and that it is not an active thermal catalyst at the temperatures that can be used for  $Co_2(CO)_8$ . We thus wondered whether the  $Co_2(CO)_8$  catalyst could be regenerated from irradiation of  $R_3 SiCo(CO)_4$  according to equations (1) and (3).

$$2 \cdot \operatorname{Co(CO)}_{4} \longrightarrow \operatorname{Co}_{2}(\operatorname{CO})_{8} \tag{3}$$

The relative importance of the two photoreactions represented by equations (1) and (2) is of fundamental interest, since there is considerable ambiguity concerning the importance of <u>primary</u> photogeneration of 17-valence electron vs. 16-valence electron fragments from coordinatively saturated, 18-valence electron species having both 2- and 1-electron donor ligands. For  $R_3SiCo(CO)_4$  the orbital diagram in Scheme I is appropriate, and it is apparent that all lowest excited states involve population of  $\sigma^*(d_2)$ 



Scheme I. One-electron orbital diagram for  $R_3SiCo(CO)_4$ .

resulting in weakening of the  $\sigma\text{-bonding}$  between Co and all of the five ligands.

#### Experimental Section

Infrared spectra were taken on a Perkin-Elmer Model 180 grating spectrophotometer in matched 1.0 mm or 0.1 mm pathlength amalgam sealed NaCl cells. Low temperature spectra were taken using a Cyrogenics Technology, Inc. Spectrim II cryocooler in conjunction with the spectrophotometer; a 1.1 mm pathlength uncompensated CaF2 cell was constructed to interface with the cold head. Electronic spectra were taken on a Varian Associates Cary Model 17 uv-vis-nir spectrophotometer, in 1.0 cm quartz or 1.3 cm Pyrex cells. Gas-liquid chromatography was performed on a Varian Associates Model 1440 or 2440 gas chromatograph equipped with flame ionization detectors and interfaced with either a Hewlett-Packard model 3370S electronic recording integrator or a Varian model A25 strip chart recorder. Hydrocarbon analyses (against internal standard n-hexane) were carried out on columns of 20% propylene carbonate on Chromasorb P (Johns-Manville), 30" x 1/8", 20°C, hydrosilation products were analyzed (against internal standard n-decane) on columns of 20% β,β'-oxydepropionitrile on Gaschrom Q (Applied Science Laboratories, 1/8" x 30', 60°C). Organic products of the photolyses of Ph<sub>3</sub>SiCo(CO)<sub>A</sub> in pentene or isooctane were separated on columns of 1.5% OV-101 on Chromosorb P (1/8" x 5", 120-250°C, programmed at 10-20°C/min.) Nuclear magnetic resonance spectra were obtained with a Hitachi-Perkin Elmer R-24B or Varian T-60 spectrometer (1H, 60 MHz) or with a Jeolco FX-90Q spectrometer (1H, 90 MHz). Mass spectra were obtained on a Varian MAT 44 spectrometer, interfaced with a Varian series 1400 gas chromatograph for GC/MS.

All air-sensitive organometallic compounds were handled under an argon atmosphere by conventional Schlenk techniques, or under an atmosphere of prepurified nitrogen in a Vacuum Atmospheres Dri-Lab glove box. Photosensitive solutions were handled under red safelights or in the dark.  $Co_2(CO)_8$  (ROC/RIC, Strem) and P(OPh) $_3$  (Strem) were used as received. PPh $_3$  (Aldrich) and

HSiPh<sub>3</sub> (Petrarch) were recrystallized from ethanol and pentane, respectively, before use. Alkenes (99%, PCR or Phillips Chemical Company) were passed through activated alumina (MC&B chromatographic grade, 80-325 mesh) prior to use and stored in amber bottles at 4°C. Triethylsilane (Petrarch) was treated similarly.  $Co_4(CO)_{12}$  was prepared by heating  $Co_2(CO)_8$  under  $N_2$  in isooctane at 80-90°C for 12 hrs, filtering off the product and recrystallizing from benzene/isooctane under  $N_2$ .  $R_3SiCo(CO)_4$  (R = Et, Ph) were prepared by the literature method of Chalk and Harrod. Solvents were reagent grade or better; isooctane was spectrophotometric grade. For work with air- or water-sensitive species solvents were distilled under argon from appropriate drying agents.

Solutions for irradiation were hermetically sealed into 13 mm o.d. pyrex ampules after at least four freeze-pump-thaw degassing cycles under high vacuum. Alternatively, the organometallic was added, in the glove box to previously degassed solvent, the mixture transferred to 13 mm o.d. pyrex tubes with ground glass joint tops and sealed with rubber septa. Irradiation samples were generally 1-4 ml in volume,  $\sim 10^{-3} \text{M}$  in metal carbonyl for catalysis and  $10^{-2} - 10^{-4} \text{M}$  for spectral studies. Light intensities were determined by ferrioxalate actinometry. Near-ultraviolet irradiation was generated by two General Electric Blacklites ( $\lambda_{\text{max}} = 355 \pm 20 \text{ nm}$ , I  $\sim 2 \times 10^{-6} \text{ ein/min}$ ) or a high-pressure mercury arc lamp (Bausch and Lomb, model SP200, 200 W) with an 18 cm water filter to remove infrared radiation. Monochromatic light was supplied by medium-pressure mercury arc lamps (Hanovia, 450 W or 550 W) filtered with Corning glass filters to isolate the desired emission (I  $\sim 10^{-7} \text{ ein/min}$  with associated merry-go-round). 254 nm irradiation was supplied by a UV-Products low-pressure mercury arc lamp.

The complexes  $R_3SiCo(CO)_4$  (R = Et, Ph) were synthesized and purified according to the literature procedures, <sup>5</sup> equation (4). These complexes are

$$Co_2(CO)_8 + 2HSiR_3 \xrightarrow{\Delta} 2R_3SiCo(CO)_4 + H_2$$
 $R = Et, Ph$ 

(4)

thermally inert at 25°C in deoxygenated hydrocarbon solutions containing 1-pentene,  $\operatorname{HSiR}_3$ ,  $\operatorname{P(OPh)}_3$  or mixtures of these reagents. However, these complexes are photosensitive upon near ultraviolet irradiation. Scheme II summarizes the photochemistry under various conditions. The data support the conclusion that dissociative loss of CO, equation (2), is the main result of photoexcitation of  $\operatorname{S_3SiCo(CO)}_4$ . However, small quantum yields for  $\operatorname{Co_2(CO)}_8$  and some  $\operatorname{Co_4(CO)}_{12}$  formation in alkane solution are found and the chemical yield of  $\operatorname{Co_2(CO)}_8$  and  $\operatorname{Co_4(CO)}_{12}$  are >90% based on  $\operatorname{R_3SiCo(CO)}_4$  consumed. In the presence of 1.0  $\operatorname{M}$  1-pentene at 25°C the quantum yield for  $\operatorname{Co_2(CO)}_8$  formation increases markedly, but at -78°C the photoreaction leads to no detectable  $\operatorname{Co_2(CO)}_8$ . At -78°C infrared spectral changes accompanying 355 nm

$$R_{3}SiCo(CO)_{3}(pentene) + CO$$

$$\uparrow hv, -78^{\circ}C$$
1-pentene
$$CO + R_{3}SiCo(CO)_{3}P(OPh)_{3} \xrightarrow{hv, 25^{\circ}C} R_{3}SiCo(CO)_{4} \xrightarrow{hv, 25^{\circ}C} R'_{3}SiH$$

$$O.1 \ \underline{M} \ P(OPh)_{3}$$

$$R = Et$$

$$hv \downarrow 1 \ \underline{M} \ 1\text{-pentene}$$

$$25^{\circ}C \downarrow \Phi_{366} = 0.23$$

$$Co_{2}(CO)_{8} + Co_{4}(CO)_{12}$$

$$+$$

$$1somers of (pentenyl)SiEt_{3}$$

$$R'_{3}SiCo(CO)_{4} + R_{3}SiH$$

$$R'_{3}SiH \rightarrow R'_{3}SiH \rightarrow R'_{3}SiH \rightarrow R'_{3}SiCo(CO)_{4} + R_{3}SiH \rightarrow R'_{3}SiCo(CO)_{4} + R_{3}SiCo(CO)_{4} + R_{3}SiCo(CO)_{4}$$

Scheme II: Photochemistry of R-SiCo(CO).

irradiation in 1.0  $\underline{\text{M}}$  1-pentene in isooctane are consistent with the formation of R<sub>3</sub>SiCo(CO)<sub>3</sub>(pentene) (R = Et,  $\nu_{\text{CO}}$  = 1966 sh, 1959 cm<sup>-1</sup>; R = Ph,  $\nu_{\text{CO}}$  = 1977 sh, 1970 cm<sup>-1</sup>). Addition of P(OPh)<sub>3</sub>/isooctane in the dark to the irradiated solution at -78°C followed by warmup to 25°C and infrared analysis reveals the presence of R<sub>3</sub>SiCo(CO)<sub>3</sub>P(OPh)<sub>3</sub>, evidencing photogeneration of a labile species at -78°C. Warmup of Et<sub>3</sub>SiCo(CO)<sub>3</sub>(pentene) - containing solutions in the absence of added P(OPh)<sub>3</sub> gives regeneration of Et<sub>3</sub>SiCo(CO)<sub>4</sub> and a significant amount of Co<sub>4</sub>(CO)<sub>12</sub>. The low temperature irradiations appear to rule out chemistry according to equation (1) as the major pathway to Co<sub>2</sub>(CO)<sub>8</sub>. At 25°C in the presence of 1-pentene, irradiation of Et<sub>3</sub>SiCo(CO)<sub>4</sub> leads to formation of isomers of (penteny1)SiEt<sub>3</sub> as the dominant Si-containing product. 8 though initial infrared spectral changes are consistent with formation of Et<sub>3</sub>SiCo(CO)<sub>3</sub>(pentene).

Consistent with dissociative loss of CO as the major primary photoreaction, we find high quartum efficiency ( $\Phi \sim 0.28$ ) for the substitution of CO by P(OPh)<sub>3</sub>. Irradiation of R<sub>3</sub>SiCo(CO)<sub>4</sub> in the presence of R'<sub>3</sub>SiH, Figure 1, results in clean formation of R'<sub>3</sub>SiCo(CO)<sub>4</sub> with good quantum yield; irradiation in the presence of 0.1 M R'<sub>3</sub>SiH and 0.3 M P(OPh)<sub>3</sub> only yields R<sub>3</sub>SiCo(CO)<sub>3</sub>(P(OPh)<sub>3</sub>). The rationale for these results is given by equations (2) and (5)-(7), indicating competitive capture of the 16 species by P(OPh)<sub>3</sub> or HSiR<sub>3</sub>.

$$R_{3}SiCo(CO)_{3} + P(OPh)_{3} \longrightarrow R_{3}SiCo(CO)_{3}(P(OPh)_{3})$$

$$R_{3}SiCo(CO)_{3} + R'_{3}SiH \longrightarrow R_{3}SiCo(CO)_{3}(H)(SiR'_{3}) \longrightarrow R'_{3}SiCo(CO)_{3} + R_{3}SiH$$

$$R'_{3}SiCo(CO)_{3} + CO \longrightarrow R'_{3}SiCo(CO)_{4}$$

$$(7)$$

With respect to catalysis, we find that irradiation of  $R_3SiCo(CO)_4$  in the presence of 1-pentene/HSiEt $_3$  does lead to the isomerization and hydrosilation of the 1-pentene, equation (8). Only traces of (pentenyl)SiEt $_3$ '

+ HSiEt<sub>3</sub> 
$$\xrightarrow{355 \text{ nm}}$$
  $(\underline{n}\text{-pentyl})\text{SiEt}_3$  + +  $(\underline{n}\text{-pentyl})\text{SiEt}_3$  +  $(\underline{n}\text{-pentyl})\text{SiEt}_3$  +  $(\underline{n}\text{-pentane})$  (8)

R = Ph,Et +  $(\underline{n}\text{-pentane})$  + (pentenyl)SiEt<sub>3</sub>(trace)

products are found. Data for the photocatalyzed reactions are given in the Table. The R<sub>3</sub>SiCo(CO)<sub>4</sub> species lead to extensive pentene isomerization prior to significant extent conversion to  $(\underline{n}$ -pentyl)SiEt<sub>3</sub>. Of the order of  $10^3$  molecules of  $(\underline{n}\text{-pentyl})$ SiEt $_3$  have been observed per molecule of  $R_3$ SiCo(CO) $_4$  initially added and the turnover rate for formation of this product is of the order of  $25 \, h^{-1}$  at the light intensity used to generate data given in the Table. Higher light intensities have yielded observed turnover rates for the reaction of ~250 h<sup>-1</sup>. Thermal activity of  $R_3SiCo(CO)_4$  at 25°C for (<u>n</u>-pentyl)SiEt<sub>3</sub> formation is low on the scale of light induced activity, but significant isomerization is detectable in thermal controls. The Table also shows data that reveal that when thermal catalytic activity of  $Co_2(CO)_8$  is over, catalytic activity can be revived by irradiation. These representative photoactivation data show an additional ~100 turnovers per Co after the catalytic activity for the  ${\rm Co_2(CO)_8}$  is over. While it is tempting to conclude that the same catalyst is generated from irradiation of  $\mathrm{Et_3SiCo(CO)_4}$  as from  $\mathrm{Co_2(CO)_8}$  in the dark, we cannot yet unequivocally make this conclusion. We do find that the  $(\underline{n}\text{-pentyl})$ SiEt $_3$  is the dominant detectable Si-containing product in both cases. Thus, it would appear that the  $Co_2(CO)_8$  activity can be prolonged by irradiation; infrared analysis of the  $\mathrm{Co_2(CO)_8/pentene/HSiEt_3}$  solution when thermal reaction has stopped shows  $\mathrm{Et_3SiCo(CO)_4}$  to account for >80% of the Co. The photoreaction of  $\text{Et}_3 \text{SiCo(CO)}_4$  in pentene to yield (pentenyl)SiEt $_3$  (vide supra) provides a pathway to an active catalyst. The formation of (pentenyl)SiEt, may occur by first forming Et<sub>3</sub>SiCo(CO)<sub>3</sub>(pentene) followed by insertion to yield (pentenylSiEt $_3$ )Co(CO) $_3$  which then rapidly gives  $\beta$ -hydrogen transfer to yield the substitution labile complex (pentenylSiEt $_3$ )Co(CO) $_3$ H. This hydride can then lose the (pentenyl)SiEt<sub>3</sub> to leave behind the coordinatively

unsaturated  $HCo(CO)_3$  that is very likely the active catalyst in the  $Co_2(CO)_8$  system.<sup>2,5</sup> In addition to determining its role in forming the active catalyst here, further study of this photoreaction is being undertaken to gain insight into the possible mechanism in  $Fe(CO)_5$  photocatalyzed reaction of  $HSiR_3/1$ -pentene that gives substantial amounts of (pentenyl)SiR<sub>3</sub> as a catalysis product.<sup>8,9</sup>

Acknowledgements. We thank the Office of Naval Research for partial support of this research and M.S.W. acknowledges support as a Dreyfus Teacher-Scholar grant recipient, 1975-1980. C.L.R. was supported during Spring, 1980 by a Fellowship with funds generously provided by Eastman Kodak Co.

#### References

- 1. Reichel, C.L.; Wrighton, M.S. J. Am. Chem. Soc., 1979, 101, 6769.
- 2. Harrod, J.F.; Chalk, A.J. "Organic Synthesis via Metal Carbonyls", Vol. 2, Pino, P.; Wender, I. eds., John Wiley & Sons, Inc.: New York, 1977; pp. 687-690.
- 3. (a) Geoffroy, G.L.; Wrighton, M.S. "Organometallic Photochemistry", Academic Press, Inc.: New York, 1979; (b) Severson, R.G.; Wojcicki, A. J. Organometal. Chem., 1978, 157, 173.
- 4. Photoelectron spectroscopy of  $H_3SiCo(CO)_4$  shows the  $3d_x2_y2$ ,  $d_{xy}$  and  $3d_{xz}$ ,  $d_{yz}$  orbitals to be the highest occupied levels with the  $\sigma_b$  level more stable: Cradock, S.; Ebsworth, E.A.V.; Robertson, A. <u>J. Chem. Soc.</u>, Dalton, 1973, 22.
- 5. Chalk, A.J.; Harrod, J.F. J. Am. Chem. Soc., 1967, 89, 1640.
- 6. Hatchard, C.G.; Parker, C.A. Proc. Roy. Soc. A, 1956, 235, 518.
- 7. These  $R_3SiCo(CO)_3P(OPh)_3$  complexes have ir spectra in the CO region as follows: R = Et, 1969 sh, 1965 cm<sup>-1</sup>; R = Ph, 1980 sh, 1973 cm<sup>-1</sup> in isooctane/1-pentene (1/1 by volume). These spectra are similar to those for the  $R_3SiCo(CO)_3$  (pentene) detected at -78°C consistent with a similar arrangement of the CO groups. The spectra accord well with those for  $R_3ECo(CO)_3L$  complexes previously reported: Boyd, T.E.; Brown, T.L. Inorg. Chem., 1974, 13, 422.
- 8. The isomers have been characterized by GC and mass spectral analyses compared to authentic samples prepared previously: Schroeder, M.A.; Wrighton, M.S. J. Organometal. Chem., 1977, 128, 345.
- 9. (a) Austin, R.G.; Paonessa, R.S.; Giordano, P.J.; Wrighton, M.S. Adv. Chem. Ser., 1978, 168, 189; (b) Nesmeyanov, A.N.; Friedlina, R.K.; Chukovskaya, E.C.; Petrova, R.G.; Belyavsky, A.B. Tetrahedron, 1962, 17, 61; (c) Friedlina, R.K.; Chukovskaya, E.C.; Tsao, J.; Nesmeyanov, A.N.; Dokl. Akad. Nauk. SSSR, 1960, 122, 37.

ದ
HSiEt <sub>3</sub> .ª
with
1-Pentene
of
Reaction
ocatalyzed
Photo
Table.

Product Analysis, %

the engineering of the first of the second o

Catalyst Precursor	1-Pentene/SiEt <sub>3</sub>	Irrdn Time	% Consumption of 1-Pentene	$\overline{n}$ -pentane	<u>t</u> -2-	<u>c</u> -2-pentene	$(\frac{n}{2}$ -pentyl) SiEt <sub>3</sub>
PhaSico(c0)	MO.1/M 0.1	0 (39h thermal control) <sup>b</sup>	8	0	16	0	6
$(1.79\times10^{-3}M)$	(in $\underline{n}$ -octane)	-	26	$\nabla$	8	7	4
		4.7	96	$\nabla$	64	16	50
•		39	26	~	22	13	30
PhaSico(co)	3.3M/3.3M	-	7.9	_	18	14	9
$(1.79\times10^{-3}M)$	(neat)	4.7	26	$\nabla$	79	14	6
		39	>97	₽	38	6	52
Et 251Co(CO) 2	3.3M/3.3M	4.7	97	~	79	14	8
$(1.79\times10^{-3}M)$	(neat)	39	>97	<b>-</b> -	32	<b>&amp;</b>	09
8 (00) <b>200</b>	3.3M/3.3M	0 (10h thermal) <sup>c</sup>	>97			1 1 5 8	23
$(2\times10^{-4})$	(near)	15 after 10h thermal <sup>u</sup>	>97			-	35
8 <sup>(00)</sup> 200	3.3M/3.3M	0 (17h thermal) <sup>C</sup>	×97			-	15
$(2\times10^{-3}\overline{M})$	(neat)	10h after 17h thermal <sup>u,e</sup>	. Ye<				28
					ľ		

<sup>a</sup>Irradiation of 1.0 cc deoxygenated samples in Pyrex ampules using 355  $\pm$  20 nm light at 2  $\times$  10<sup>-6</sup> ein/min and at 25°C. Solutions were analyzed by GC and quantitated by using internal standards n-hexane for determination of C<sub>5</sub>'s and n-decane for (pentyl)SiEt<sub>3</sub>.


Note:  $\frac{n}{3}$  h under same conditions except not irradiated.

This represents a point in the reaction when there is no further detectable activity.

 $^{
m d}_{
m In}$  these experiments the irradiation was not started until thermal activity ceased. The species irradiated is principally Et $_3$ SiCo(CO) $_4$  as detected by infrared.

eA higher light intensity was used in this experiment.

Figure Caption. Comparison of photochemistry of  $Ph_3SiCo(CO)_4$  in the presence of  $HSiEt_3$  only (a) and  $HSiEt_3$  and  $P(OPh)_3$  (b). Irradiation of the  $25^{\circ}C$  deoxygenated solutions is at 355 nm ( $\pm$  20 nm). Note that quantum yield for disappearance of  $Ph_3SiCo(CO)_4$  is the same in both cases but that silane exchange occurs exclusively in (a) and  $P(OPh)_3$  substitution occurs exclusively in (b), consistent with competitive capture of a  $16e^-$  species by  $P(OPh)_3$  or  $HSiEt_3$ , equations (5) and (6) in text.



### TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies		No. Copies
Office of Naval Research		U.S. Army Research Office	
Attn: Code 472		Attn: CRD-AA-IP	
800 North Quincy Street		P.O. Box 1211	
Arlington, Virginia 22217	2	Research Triangle Park, N.C. 27709	1
ONR Branch Office		Naval Ocean Systems Center	
Attn: Dr. George Sandoz		Attn: Mr. Joe McCartney	
536 S. Clark Street		San Diego, California 92152	1
Chicago, Illinois 60605	1		
		Naval Weapons Center	
ONR Area Office		Attn: Dr. A. B. Amster,	
Attn: Scientific Dept.		Chemistry Division	•
715 Broadway	•	China Lake, California 93555	1
New York, New York 10003	1	Name 1 Civil Produceming Laboratory	
OND II and a Dead of 1 Offi		Naval Civil Engineering Laboratory	
ONR Western Regional Office		Attn: Dr. R. W. Drisko	,
1030 East Green Street	•	Port Hueneme, California 93401	1
Pasadena, California 91106	1	Department of Physics Chemistry	
OND Distance (Control Doods and Office		Department of Physics & Chemistry	
ONR Eastern/Central Regional Office		Naval Postgraduate School Monterey, California 93940	1
Attn: Dr. L. H. Peebles		Monterey, Carronnia 95940	•
Building 114, Section D 666 Summer Street		Dr. A. L. Slafkosky	
Boston, Massachusetts 02210	1	Scientific Advisor	
boston, lassachusetts 02210	•	Commandant of the Marine Corps	
Director, Naval Research Laboratory		(Code RD-1)	_
Attn: Code 6100		Washington, D.C. 20380	1
Washington, D.C. 20390	1	000	
		Office of Naval Research	
The Assistant Secretary		Attn: Dr. Richard S. Miller	
of the Navy (RE&S)		800 N. Quincy Street	,
Department of the Navy		Arlington, Virginia 22217	1
Room 4E736, Pentagon	,	Naval Chin Research and Development	
Washington, D.C. 20350	1	Naval Ship Research and Development Center	
Commander, Naval Air Systems Command	l	Attn: Dr. G. Bosmajian, Applied	
Attn: Code 310C (H. Rosenwasser)		Chemistry Division	_
Department of the Navy		Annapolis, Maryland 21401	1
Washington, D.C. 20360	1	Naval Ocean Systems Center	
Defense Technical Information Center	•	Attn: Dr. S. Yamamoto, Marine	
Building 5, Cameron Station		Sciences Division	
Alexandria, Virginia 22314	12	San Diego, California 91232	1
Dr. Fred Saalfeld		Mr. John Boyle	
Chemistry Division, Code 6100		Materials Branch	
Naval Research Laboratory	_	Naval Ship Engineering Center	•
Washington, D.C. 20375	1	Philadelphia, Pennsylvania 19112	1

SP472-3/A3

472:GAN:716:ddc 78u472-608

# TECHNICAL REPORT DISTRIBUTION LIST, GEN

	No. Copies
Dr. Rudolph J. Marcus Office of Naval Research Scientific Liaison Group American Embassy APO San Francisco 96503	1
Mr. James Kelley DTNSRDC Code 2803 Annapolis, Maryland 21402	1

#### TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. Paul Delahay		Dr. P. J. Hendra	
Department of Chemistry		Department of Chemistry	
New York University		University of Southhampton	
New York, New York 10003	1	Southhampton SO9 5NH United Kingdom	1
Dr. E. Yeager		Chiled Kingdom	•
Department of Chemistry		Dr. Sam Perone	
Case Western Reserve University		Department of Chemistry	
Cleveland, Ohio 41106	1	Purdue University	
·		West Lafayette, Indiana 47907	1
Dr. D. N. Bennion		• •	
Department of Chemical Engineering		Dr. Royce W. Murray	
Brigham Young University		Department of Chemistry	
Provo, Utah 84602	1	University of North Carolina Chapel Fill, North Carolina 27514	1
Dr. R. A. Marcus		onaper will, worth darolina 2/314	•
Department of Chemistry		Naval Ocean Systems Center	
California Institute of Technology		Attn: Technical Library	
Pasadena, California 91125	1	San Diego, California 92152	1
Dr. J. J. Auborn		Dr. C. F. Mueller	
Bell Laboratories		The Electrochemistry Branch	
Murray Hill, New Jersey 07974	1	Materials Division, Research & Technology Department	
Dr. Adam Heller		Naval Surface Weapons Center	
Bell Laboratories		White Oak Laboratory	
Murray Hill, New Jersey 07974	1	Silver Spring, Marvland 20910	1
Dr. T. Katan		Dr. G. Goodman	
Lockheed Missiles & Space		Globe-Union Incorporated	
Co, Inc.		5757 North Green Bay Avenue	
P.O. Box 504		Milwaukee, Wisconsin 53201	1
Sunnyvale, California 94088	1	•	
		Dr. J. Boechler	
Dr. Joseph Singer, Code 302-1		Electrochimica Corporation	
NASA-Lewis		Attention: Technical Library	
21000 Brookpark Road	_	2485 Charleston Road	
Cleveland, Ohio 44135	1	Mountain View, California 94040	1
Dr. B. Brummer		Dr. P. P. Schmidt	
EIC Incorporated		Department of Chemistry	
55 Chapel Street		Oakland University	
Newton, Massachusetts 02158	1	Rochester, Michigan 48063	1
Library		Dr. H. Richtol	
P. R. Mallory and Company, Inc.		Chemistry Department	
Northwest Industrial Park		Rensselaer Polytechnic Institute	
Burlington, Massachusetts 01803	1	Troy, New York 12181	1

#### TECHNICAL REPORT DISTRIBUTION LIST, 359

	No. Copies		No. Copies
Dr. A. B. Ellis		Dr. R. P. Van Duyne	
Chemistry Department		Department of Chemistry	
University of Wisconsin	_	Northwestern University	١ .
Madison, Wisconsin 53706	1	Evanston, Illinois 60201	' 1
Dr. M. Wrighton		Dr. B. Stanley Pons	
Chemistry Department		Department of Chemistry	
Massach seres Institute		University of Alberta	
of Technology		Edmonton, Alberta	
cambridge, Massachusetts 02139	1	CANADA T6G 2G2	1
Larry E. Plew		Dr. Michael J. Weaver	
Naval Weapons Support Center		Department of Chemistry	
Code 30736, Building 2906		Michigan State University	
Crane, Indiana 47522	1	East Lansing, Michigan 48824	1
S. Ruby		Dr. R. David Rauh	
DOE (STOR)		EIC Corporation	
600 E Street		55 Chapel Street	
Washington, D.C. 20545	1	Newton, Massachusetts 02158	1
<b>D</b> 4			
Dr. Aaron Wold		Dr. J. David Margerum	
Brown University		Research Laboratories Division	
Department of Chemistry	_	Hughes Aircraft Company	
Providence, Rhode Island 02192	1	3011 Malibu Canyon Road	_
		Malibu, California 90265	I
Dr. R. C. Chudacek			
McGraw-Edison Company		Dr. Martin Fleischmann	
Edison Battery Division		Department of Chemistry	
Post Office Box 28	•	University of Southampton	
Bloomfield, New Jersey 07003	1	Southampton 509 5NH England	1
Dr. A. J. Bard		Dr. Janet Osteryoung	
University of Texas		Department of Chemistry	
Department of Chemistry		State University of New	
Austin, Texas 78712	1	York at Buffalo	
		Buffalo, New York 14214	1
Dr. M. M. Nicholson			
Electronics Research Center		Dr. R. A. Osteryoung	
Rockwell International		Department of Chemistry	
3370 Miraloma Avenue		State University of New	
Anaheim, California	1	York at Buffalo	
		Buffalo, New York 14214	1
Dr. Donald W. Ernst			
Naval Surface Weapons Center		Mr. James R. Moden	
Code R-33		Naval Underwater Systems	
White Oak Laboratory		Center	
Silver oring, Maryland 20910	1	Code 3632	
		Newport, Rhode Island 02840	1

## TECHNICAL REPORT DISTRIBUTION LIST, 359

	No.		No.
	Copies		Copies
Dr. R. Nowak		Dr. John Kincaid	1
Naval Research Laboratory		Department of the Navy	
Code 6130		Stategic Systems Project Office	
Washington, D.C. 20375	1	Room 901	
		Washington, DC 20376	
Dr. John F. Houlihan		,	
Shenango Valley Campus		M. L. Robertson	
Pennsylvania State University		Manager, Electrochemical	
Sharon, Pennsylvania 16146	1	Power Sonices Division	
•		Naval Weapons Support Center	
Dr. M. G. Sceats		Crane, Indiana 47522	1
Department of Chemistry			
University of Rochester		Dr. Elton Cairns	
Rochester, New York 14627	I	Energy & Environment Division	
·		Lawrence Berkeley Laboratory	
Dr. D. F. Shriver		University of California	
Department of Chemistry		Berkeley, California 94720	1
Northwestern University			
Evanston, Illinois 60201	1	Dr. Bernard Spielvogel	
•		U.S. Army Research Office	
Dr. D. H. Whitmore		P.O. Box 12211	
Department of Materials Science		Research Triangle Park, NC 27709	1
Northwestern University		•	
Evanston, Illinois 60201	1	Dr. Denton Elliott	
•		Air Force Office of	
Dr. Alan Bewick		Scientific Research	
Department of Chemistry		Bldg. 104	
The University		Bolling AFB	
Southampton, SO9 5NH England	1	Washington, DC 20332	1
Dr. A. Himy			
NAVSEA-5433	•		
NC #4			
2541 Jefferson Davis Highway			
Arlington, Virginia 20362	1		

# TECHNICAL REPORT DISTRIBUTION LIST, 051A

	No. Copies		No. Copies
Dr. M. A. El-Sayed Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1	Dr. M. Rauhut Chemical Research Division American Cyanamid Company Bound Brook, New Jersey 08805	1
Dr. E. R. Bernstein Department of Chemistry Colorado State University	-	Dr. J. I. Zink Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1
Fort Collins, Colorado 80521	1	200	•
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. D. Haarer IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. J. R. MacDonald Chemistry Division Naval Research Laboratory Code 6110	,	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Washington, D.C. 20375  Dr. G. B. Schuster Chemistry Department University of Illinois	1	Dr. William M. Jackson Department of Chemistry Howard University Washington, DC 20059	1
Urbana, Illinois 61801  Dr. A. Adamson Department of Chemistry University of Southern	ī	Dr. George E. Walraffen Department of Chemistry Howard University Washington, DC 20059	1
California Los Angeles, California 90007	1		
Dr. M. S. Wrighton Department of Chemistry Massachusett Institute of			
Technology Caroridge, Massachusetts 02139	1		